

Spin chains and combinatorics: twisted boundary conditions

A. V. Razumov, Yu. G. Stroganov

*Institute for High Energy Physics
142280 Protvino, Moscow region, Russia*

Abstract

The finite XXZ Heisenberg spin chain with twisted boundary conditions was considered. For the case of even number of sites N , anisotropy parameter $-1/2$ and twisting angle $2\pi/3$ the Hamiltonian of the system possesses an eigenvalue $-3N/2$. The explicit form of the corresponding eigenvector was found for $N \leq 12$. Conjecturing that this vector is the ground state of the system we made and verified several conjectures related to the norm of the ground state vector, its component with maximal absolute value and some correlation functions, which have combinatorial nature. In particular, the squared norm of the ground state vector is probably coincides with the number of half-turn symmetric alternating sign $N \times N$ matrices.

The Hamiltonian of the periodic XXZ Heisenberg spin chain with the number of sites N and the anisotropy parameter Δ has the form

$$H^{(N)} = - \sum_{i=1}^N [\sigma_i^x \sigma_{i+1}^x + \sigma_i^y \sigma_{i+1}^y + \Delta \sigma_i^z \sigma_{i+1}^z].$$

where we assume that

$$\sigma_{N+1}^x = \sigma_1^x, \quad \sigma_{N+1}^y = \sigma_1^y, \quad \sigma_{N+1}^z = \sigma_1^z.$$

In terms of the operators $\sigma_i^+ = (\sigma_i^x + i\sigma_i^y)/2$, $\sigma_i^- = (\sigma_i^x - i\sigma_i^y)/2$ and σ_i^z one has the following expression for the Hamiltonian

$$H^{(N)} = -2 \sum_{i=1}^N [\sigma_i^+ \sigma_{i+1}^- + \sigma_i^- \sigma_{i+1}^+] - \Delta \sum_{i=1}^N \sigma_i^z \sigma_{i+1}^z, \quad (1)$$

and the boundary conditions take the form

$$\sigma_{N+1}^+ = \sigma_1^+, \quad \sigma_{N+1}^- = \sigma_1^-, \quad \sigma_{N+1}^z = \sigma_1^z. \quad (2)$$

It was proved in paper [1] that for $\Delta = -1/2$ and for an odd N the Hamiltonian $H^{(N)}$ has the eigenvalue $-3N/2$, see also [2]. In our paper [3] we analysed the explicit expressions for the corresponding eigenvectors and made the conjecture, which was verified for $N \leq 17$, that the eigenvalue $-3N/2$ corresponds to the ground state of the

system. This is in agreement with the earlier numerical results reported in paper [4]. In the same paper [3] we made and verified for $N \leq 17$ a lot of conjectures concerning the norm of the ground state, the largest components of the ground state vector and some correlation functions. These conjectures have a combinatorial character and are related, in particular, to the number of alternating sign matrices (see, for example, [5, 6]).

Several variations of the XXZ Heisenberg chain with a special ground state energy are also known [4, 7, 8, 9]. Very recently Batchelor, de Gier and Nienhuis considered these variations along with the corresponding $O(n)$ loop model at $n = 1$ [10]. They found the generalization for two of our conjectures for these systems and extended a list of related combinatorial objects.

First of all the authors of paper [10] reported about XXZ Heisenberg chain with twisted boundary conditions. The Hamiltonian in this case has again form (1), but instead of (2) one uses the boundary conditions

$$\sigma_{N+1}^+ = e^{i\phi} \sigma_1^+, \quad \sigma_{N+1}^- = e^{-i\phi} \sigma_1^-, \quad \sigma_{N+1}^z = \sigma_1^z.$$

Explicitly one can write

$$H^{(N)} = -2 \sum_{i=1}^{N-1} [\sigma_i^+ \sigma_{i+1}^- + \sigma_i^- \sigma_{i+1}^+] - 2 e^{-i\varphi} \sigma_N^+ \sigma_1^- - 2 e^{i\varphi} \sigma_N^- \sigma_1^+ - \Delta \sum_{i=1}^N \sigma_i^z \sigma_{i+1}^z. \quad (3)$$

In paper [4] on the base of numerical calculations it was conjectured that for the special value of the twisting angle $\phi = 2\pi/3$ and $\Delta = -1/2$ the ground state energy is equal to $-3N/2$. The proof of the existence of such an eigenvalue of the Hamiltonian was given in paper [9]. It was the case that was considered by the authors of paper [10]. They formulated two conjectures concerning the sum of the components of the supposed ground state vector and the sum of its squared components enjoying remarkable combinatorial character.

In the present paper we also consider the case $\phi = 2\pi/3$, $\Delta = -1/2$ and study the eigenvectors of the Hamiltonian (3) corresponding to the eigenvalue $-3N/2$ with the main purpose to find conjectures on correlation functions. We start with exposition of the explicit form of the eigenvectors for $N = 2, 4, 6$. We hope that this information will help someone for further treating of the problem. As a matter of fact we found explicitly the eigenvectors for $N = 8, 10$ and 12 as well. But they have 70, 252 and 924 components, respectively, so we decided against their exposition. Let us fix the notations.

We use the basis in the state space formed by the normed common eigenvectors of the operators σ_i^z . Here we associate $\uparrow$ with the eigenvalue $+1$ and $\downarrow$ with the eigenvalue -1 . Thus, the basis vectors are marked by words consisting of N letters $\uparrow$ and $\downarrow$. The set of such words is denoted $W^{(N)}$ and we write for a general state vector

$$|\Psi\rangle = \sum_{A \in W^{(N)}} \Psi_A |A\rangle.$$

Furthermore, we use such representation for the operators σ_i^x and σ_i^y that their matrix elements coincide with usual Pauli matrices. Finally, the eigenvector of the Hamiltonian (3) with the eigenvalue $-3N/2$ is denoted $|\Psi^{(N)}\rangle$.

The eigenvector $|\Psi^{(2)}\rangle$ has the form

$$|\Psi^{(2)}\rangle = -\frac{i}{2}(1 + i\sqrt{3})|\downarrow\uparrow\rangle + \frac{i}{2}(1 - i\sqrt{3})|\uparrow\downarrow\rangle.$$

Let us discuss the used normalization condition. Consider the operator

$$R = \sigma_1^x \cdots \sigma_N^x$$

which reverse the z -action projection of the spins and the antiunitary operator K of complex conjugation acting on a general state vector as

$$K|\Psi\rangle = \sum_A \Psi_A^* |A\rangle.$$

It is clear that the Hamiltonian (3) commutes with the operator $R' = RK^1$. Moreover, we have $R'^2 = 1$. We have found that for even $N \leq 12$ the eigenvalue $-3N/2$ is non-degenerate. It follows from these facts that multiplying $|\Psi^{(N)}\rangle$ by an appropriate phase factor one can satisfy the condition

$$R'|\Psi^{(N)}\rangle = |\Psi^{(N)}\rangle. \quad (4)$$

This condition is invariant under the multiplication of $|\Psi^{(N)}\rangle$ by a real number. To fix this freedom we divide $|\Psi^{(N)}\rangle$ by the least absolute value of its components. After this the vector $|\Psi^{(N)}\rangle$ is defined up to a sign which is fixed by the requirement that the sum of the components of $|\Psi^{(N)}\rangle$ be positive².

Under the above normalization conditions the vector $|\Psi^{(4)}\rangle$ looks as

$$\begin{aligned} |\Psi^{(4)}\rangle = & \frac{1}{2}(1 - i\sqrt{3})|\downarrow\downarrow\uparrow\uparrow\rangle + \frac{1}{2}(3 - i\sqrt{3})|\downarrow\uparrow\downarrow\uparrow\rangle + |\downarrow\uparrow\uparrow\downarrow\rangle \\ & + |\uparrow\downarrow\downarrow\uparrow\rangle + \frac{1}{2}(3 + i\sqrt{3})|\uparrow\downarrow\uparrow\downarrow\rangle + \frac{1}{2}(1 + i\sqrt{3})|\uparrow\uparrow\downarrow\downarrow\rangle. \end{aligned}$$

For the case $N = 6$ we obtain

$$\begin{aligned} |\Psi^{(6)}\rangle = & -i|\downarrow\downarrow\downarrow\uparrow\uparrow\uparrow\rangle + \frac{1}{2}(\sqrt{3} - 5i)|\downarrow\downarrow\uparrow\downarrow\uparrow\uparrow\rangle \\ & + (\sqrt{3} - 2i)|\downarrow\downarrow\uparrow\uparrow\downarrow\uparrow\rangle + \frac{1}{2}(\sqrt{3} - i)|\downarrow\downarrow\uparrow\uparrow\uparrow\downarrow\rangle + (\sqrt{3} - 2i)|\downarrow\uparrow\downarrow\downarrow\uparrow\uparrow\rangle \\ & + \frac{5}{2}(\sqrt{3} - i)|\downarrow\uparrow\downarrow\uparrow\downarrow\uparrow\rangle + \frac{1}{2}(3\sqrt{3} - i)|\downarrow\uparrow\downarrow\uparrow\uparrow\downarrow\rangle + \frac{1}{2}(3\sqrt{3} - i)|\downarrow\uparrow\uparrow\downarrow\downarrow\uparrow\rangle \\ & + \frac{1}{2}(3\sqrt{3} + i)|\downarrow\uparrow\uparrow\downarrow\uparrow\downarrow\rangle + \frac{1}{2}(\sqrt{3} + i)|\downarrow\uparrow\uparrow\uparrow\downarrow\downarrow\rangle + \frac{1}{2}(\sqrt{3} - i)|\uparrow\downarrow\downarrow\downarrow\uparrow\uparrow\rangle \\ & + \frac{1}{2}(3\sqrt{3} - i)|\uparrow\downarrow\downarrow\uparrow\downarrow\uparrow\rangle + \frac{1}{2}(3\sqrt{3} + i)|\uparrow\downarrow\downarrow\uparrow\uparrow\downarrow\rangle + \frac{1}{2}(3\sqrt{3} + i)|\uparrow\downarrow\uparrow\downarrow\downarrow\uparrow\rangle \\ & + \frac{5}{2}(\sqrt{3} + i)|\uparrow\downarrow\uparrow\downarrow\uparrow\downarrow\rangle + (\sqrt{3} + 2i)|\uparrow\downarrow\uparrow\uparrow\downarrow\downarrow\rangle + \frac{1}{2}(\sqrt{3} + i)|\uparrow\uparrow\downarrow\downarrow\downarrow\uparrow\rangle \\ & + (\sqrt{3} + 2i)|\uparrow\uparrow\downarrow\downarrow\uparrow\downarrow\rangle + \frac{1}{2}(\sqrt{3} + 5i)|\uparrow\uparrow\downarrow\uparrow\downarrow\downarrow\rangle + i|\uparrow\uparrow\uparrow\downarrow\downarrow\downarrow\rangle. \end{aligned}$$

¹The Hamiltonian (3) also commutes with the product of the operator reversing the direction of the chain and the operator K and such symmetry can be also used for the fixation of $|\Psi^{(N)}\rangle$.

²Note that this sum is real due to the condition (4).

Due to the complex character of the components of the eigenvector $|\Psi^{(N)}\rangle$, our data is less than for the periodic case [3]. Nevertheless, we are able to formulate and verify analogues of conjectures given in paper [3]. The first conjecture associates the eigenvalue $-3N/2$ with the ground state.

Conjecture 1 *The ground state of Hamiltonian (3) for an even N , $\phi = 2\pi/3$ and $\Delta = -1/2$ has the energy $E = -3N/2$ and the total spin z -axes projection $S_z = 0$.*

Consider now the maximal and the minimal absolute value of the components of $|\Psi^{(N)}\rangle$. Note that the minimal absolute value is taken, in particular, by the ferromagnetic component $\Psi_{\downarrow\ldots\downarrow\uparrow\ldots\uparrow}^{(N)}$ and the maximal absolute value is taken by the antiferromagnetic component $\Psi_{\downarrow\uparrow\ldots\downarrow\uparrow}^{(N)}$. For the absolute value of the ratio $\Psi_{\downarrow\uparrow\ldots\downarrow\uparrow}^{(2n)}/\Psi_{\downarrow\ldots\downarrow\uparrow\ldots\uparrow}^{(2n)}$ we have the sequence

$$1, \quad \sqrt{3}, \quad 5, \quad 14\sqrt{3}, \quad 198, \quad 1573\sqrt{3}, \quad \dots \quad (5)$$

Remind that for the periodic boundary conditions and an odd number of sites we have obtained [3] that the similar ratio $\Psi_{\downarrow\uparrow\ldots\downarrow\uparrow}^{(2n+1)}/\Psi_{\downarrow\downarrow\ldots\downarrow\uparrow\ldots\uparrow}^{(2n+1)}$ coincides with the number A_n of the alternating sign $n \times n$ matrices. The numbers A_n are given by the formula

$$A_n = \prod_{i=0}^{n-1} \frac{(3i+1)!}{(n+i)!}.$$

and the sequence A_n goes as

$$1, \quad 2, \quad 7, \quad 42, \quad 429, \quad 7436, \quad \dots \quad (6)$$

Dividing sequence (5) by sequence (6), we come to

$$1, \quad \frac{\sqrt{3}}{2}, \quad \frac{5}{7}, \quad \frac{\sqrt{3}}{3}, \quad \frac{6}{13}, \quad \frac{11\sqrt{3}}{4 \cdot 13}, \quad \dots \quad (7)$$

Let us now we pay the special attention to the largest prime divisors in the denominators, 7 for $n = 3$ and 13 for $n = 5$. Their locations prompt to consider the auxiliary sequence

$$1, \quad 4, \quad 4 \cdot 7, \quad 4 \cdot 7 \cdot 10, \quad 4 \cdot 7 \cdot 10 \cdot 13, \quad 4 \cdot 7 \cdot 10 \cdot 13 \cdot 16, \quad \dots \quad (8)$$

and to multiply sequence (7) by sequence (8). This leads to

$$1, \quad 2\sqrt{3}, \quad 2^2 \cdot 5, \quad \frac{2^3 \cdot 5 \cdot 7\sqrt{3}}{3}, \quad 2^4 \cdot 3 \cdot 5 \cdot 7, \quad 2^5 \cdot 5 \cdot 7 \cdot 11\sqrt{3}, \quad \dots \quad (9)$$

The largest prime divisors in the numerators hint at another auxiliary sequence

$$1, \quad 3, \quad 3 \cdot 5, \quad 3 \cdot 5 \cdot 7, \quad 3 \cdot 5 \cdot 7 \cdot 9, \quad 3 \cdot 5 \cdot 7 \cdot 9 \cdot 11, \quad \dots \quad (10)$$

Dividing sequence (9) by sequence (10), we arrive at

$$1, \quad \frac{2}{\sqrt{3}}, \quad \left(\frac{2}{\sqrt{3}}\right)^2, \quad \left(\frac{2}{\sqrt{3}}\right)^3, \quad \left(\frac{2}{\sqrt{3}}\right)^4, \quad \left(\frac{2}{\sqrt{3}}\right)^5, \quad \dots$$

As the result we obtain

Conjecture 2 *The absolute value of the ratio of a component of $|\Psi^{(2n)}\rangle$ with the largest absolute value to the component with the smallest absolute value is given by the formula*

$$\left| \Psi_{\downarrow\uparrow\ldots\downarrow\uparrow}^{(2n)} / \Psi_{\downarrow\ldots\downarrow\uparrow\ldots\uparrow}^{(2n)} \right| = \left(\frac{2}{\sqrt{3}} \right)^{n-1} \frac{1 \cdot 3 \cdot 5 \cdots (2n-3)(2n-1)}{1 \cdot 4 \cdot 7 \cdots (3n-5)(3n-2)} A_n.$$

Consider now the squared norm of the vector $|\Psi^{(N)}\rangle$:

$$\mathcal{N}_N^2 = \langle \Psi^{(N)} | \Psi^{(N)} \rangle = \sum_{A \in W^{(N)}} |\Psi_A^{(N)}|^2.$$

Fixing the normalization in the way discussed above we see that the sequence formed by the numbers $\mathcal{N}_{2n}^2$ goes as

$$2, \quad 10, \quad 140, \quad 5544, \quad 622908, \quad 198846076, \quad \dots \quad (11)$$

For the case of periodic boundary conditions the corresponding sequence of squared norms $\mathcal{N}_{2n+1}^2$ is

$$1, \quad 3, \quad 25, \quad 588, \quad 39204, \quad 7422987, \quad \dots \quad (12)$$

and we conjectured in paper [3] that

$$\mathcal{N}_{2n+1}^2 = \left(\frac{3}{4} \right)^n \left[\frac{2 \cdot 5 \cdots (3n-1)}{1 \cdot 3 \cdots (2n-1)} \right]^2 A_n^2.$$

Dividing sequence (11) by the sequence formed by the numbers A_n^2 , we obtain

$$2, \quad \frac{5}{2}, \quad \frac{2^2 \cdot 5}{7}, \quad \frac{2 \cdot 11}{7}, \quad \frac{2^2 \cdot 11}{13}, \quad \frac{11 \cdot 17}{2^2 \cdot 13}, \quad \dots$$

Using the same method as for conjecture 2, we come to the following analogue of conjecture 3 formulated in paper [3].

Conjecture 3 *The squared norm of the vector $|\Psi^{(2n)}\rangle$ is given by the formula*

$$\mathcal{N}_{2n}^2 = \frac{2 \cdot 5 \cdots (3n-1)}{1 \cdot 4 \cdots (3n-2)} A_n^2.$$

Let us remark that joining the sequences (11) and (12) we come to the sequence

$$1, \quad 2, \quad 3, \quad 10, \quad 25, \quad 140, \quad 588, \quad 5544, \quad 39204, \quad 622908, \quad 7422987, \quad \dots$$

This is the sequence formed by the number of half-turn symmetric alternating sign $N \times N$ matrices A_N^{HT} (see [11, 12] and references therein). Therefore we can combine our conjecture 3 and conjecture 3 of paper [3] into the statement that $\mathcal{N}_N^2 = A_N^{HT}$.

Batchelor, de Gier and Nienhuis discovered another analogue of the conjecture 3 of paper [3].

Conjecture 3' (Batchelor, de Gier and Nienhuis [10]) *The sum of the squared components of the vector $|\Psi^{(2n)}\rangle$ is given by*

$$\sum_{A \in W^{(2n)}} [\Psi_A^{(2n)}]^2 = A_n^2.$$

The same authors gave the analogue of conjecture 4 from paper [3].

Conjecture 4 (Batchelor, de Gier and Nienhuis [10]) *The sum of the components of the vector $|\Psi^{(2n)}\rangle$ is given by*

$$\sum_{A \in W^{(2n)}} \Psi_A^{(2n)} = 3^{n/2} A_n.$$

Note that Hamiltonian (3) has a modified shift invariance. Indeed, it commutes with the operator

$$T' = T \exp \left[\frac{i\phi}{2} \left(\sigma_N^z - \frac{1}{N} \sum_{i=1}^N \sigma_i^z \right) \right],$$

where T is the operator of the right shift by a site. Our choice of the operator of modified shift invariance is dictated by the requirement $T'^N = 1$. Note that the operator T' commutes with the operator R' . It follows from these facts and from requirement (4) that in the case under consideration either $T'|\Psi^{(N)}\rangle = |\Psi^{(N)}\rangle$ or $T'|\Psi^{(N)}\rangle = -|\Psi^{(N)}\rangle$. It appears that for even $N \leq 12$ one has

$$T'|\Psi^{(N)}\rangle = |\Psi^{(N)}\rangle. \quad (13)$$

It is natural to suppose that it is so for any even N . It can be deduced from (13) that for any $k = 0, 1, \dots$

$$\sum_{A \in W^{(N)}} [\Psi_A^{(N)}]^{3(2k+1)} = 0.$$

Let us proceed to the consideration of correlation functions. For any operator O we denote

$$\langle O \rangle_N = \frac{\langle \Psi^{(N)} | O | \Psi^{(N)} \rangle}{\langle \Psi^{(N)} | \Psi^{(N)} \rangle}.$$

It is evident that $\langle \sigma_{i_1}^z \dots \sigma_{i_k}^z \rangle_{2n}$ does not depend on the order of the operators σ_i^z . From (4) we obtain

$$\langle \sigma_{i_1}^z \dots \sigma_{i_{2k+1}}^z \rangle_{2n} = 0$$

for any $k = 0, 1, \dots$, and the equality (13) implies that

$$\langle \sigma_{i_1}^z \dots \sigma_{i_k}^z \rangle_{2n} = \langle \sigma_{i_1+1}^z \dots \sigma_{i_k+1}^z \rangle_{2n}.$$

To formulate our conjectures on correlation functions introduce the operators

$$\alpha_i = (1 + \sigma_i^z)/2.$$

The analogue of conjecture 5 of paper [3] is

Conjecture 5

$$\langle \alpha_i \alpha_{i+2} \rangle_{2n} = \frac{71n^4 - 19n^2 + 20}{16(4n^2 - 1)^2}.$$

Unlike the periodic case, this conjecture is a simple consequence of

Conjecture 6 *There is a simple formula for the correlation functions called the Probabilities of Formation of Ferromagnetic String [13]:*

$$\frac{\langle \alpha_1 \alpha_2 \dots \alpha_{k-1} \rangle_{2n}}{\langle \alpha_1 \alpha_2 \dots \alpha_k \rangle_{2n}} = \frac{(2k-2)!(2k-1)!(2n+k-1)!(n-k)!}{(k-1)!(3k-2)!(2n-k)!(n+k-1)!}.$$

Note that the conjectured formulas for correlation functions have the same thermodynamic limit both for the twisted and periodic boundary conditions [3].

Acknowledgments The work was supported in part by the Russian Foundation for Basic Research.

References

- [1] Yu. G. Stroganov, *The importance of being odd*, `cond-mat/0012035`
- [2] R. J. Baxter, *Adv. Stud. in Pure Math.* **19** (1989) 95
- [3] A. V. Razumov and Yu. G. Stroganov, *Spin chains and combinatorics*, `cond-mat/0012141`
- [4] F. C. Alcaraz, M. N. Barber and M. T. Batchelor, *Ann. Phys. (NY)* **182** (1988) 280
- [5] D. Bressoud, *Proofs and Confirmations*, Mathematical Association of America, Washington, DC, 1999
- [6] D. Bressoud and J. Propp, *Notices of the AMS* **46** (1999) 637
- [7] F. C. Alcaraz, M. N. Barber, M. T. Batchelor, R. J. Baxter and G. R. W. Quispel, *J. Phys.* **A20** (1987) 6397
- [8] V. Fridkin, Yu. G. Stroganov and Don Zagier, *J. Phys.* **A33** (2000) L121
- [9] V. Fridkin, Yu. G. Stroganov and Don Zagier, *J. of Stat. Phys.* **102** (2001) 781
- [10] M. T. Batchelor, J. de Gier and B. Nienhuis, *The quantum symmetric XXZ chain at $\Delta = -\frac{1}{2}$, alternating sign matrices and plane partitions*, `cond-mat/0101385`
- [11] D. P. Robbins, *Symmetry classes of alternating sign matrices*, `math.CO/0008045`
- [12] B. Kuperberg, *Symmetry classes of alternating-sign matrices under one roof*, `math.CO/0008184`
- [13] V. E. Korepin, A. G. Izergin and N. M. Bogoliubov, *Quantum Inverse Scattering Method, Correlation Functions and Algebraic Bethe Ansatz*, Cambridge University Press, 1993